


PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 209/20, A61K 31/40, C07C 311/29, 311/19		A1	(11) International Publication Number: WO 98/17645
			(43) International Publication Date: 30 April 1998 (30.04.98)
(21) International Application Number: PCT/US97/18235 (22) International Filing Date: 20 October 1997 (20.10.97) (30) Priority Data: 60/029,585 22 October 1996 (22.10.96) US (71) Applicant (for all designated States except US): PHARMACIA & UPJOHN COMPANY [US/US]; 301 Henrietta Street, Kalamazoo, MI 49001 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): WARPEHOSKI, Martha, A. [US/US]; 7600 Curry Lane, Portage, MI 49024 (US). MITCHELL, Mark, A. [US/US]; 1628 Dover Road, Kalamazoo, MI 49008 (US). JACOBSEN, E., Jon [US/US]; 74 South Lake Doster Drive, Plainwell, MI 49080 (US). (74) Agent: YANG, Lucy, X.; Pharmacia & Upjohn Company, Intellectual Property Legal Services, 301 Henrietta Street, Kalamazoo, MI 49001 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(54) Title: α -AMINO SULFONYL HYDROXAMIC ACIDS AS MATRIX METALLOPROTEINASE INHIBITORS			
(57) Abstract A compound of formula (I) or pharmaceutical acceptable salts thereof, wherein R ₁ is isopropyl, 2-methylbut-2-yl, phenyl, benzyl, or 1H-indol-3-ylmethyl; R ₂ is n-octyl, phenyl, or phenyl substituted with methoxy, fluoro, or bromo, are matrix metalloproteinase inhibitors.			



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

α-AMINO SULFONYL HYDROXAMIC ACIDS AS MATRIX METALLOPROTEINASE INHIBITORS

FIELD OF THE INVENTION

5 The present invention relates to therapeutically active α -amino sulfonyl hydroxamic acids, to pharmaceutical compositions containing them, and to the method of using such compounds. Particularly, the compounds of the invention are inhibitors of matrix metalloproteinases involved in tissue degradation.

10 BACKGROUND OF THE INVENTION

Loss of connective tissue integrity occurs in many disease processes, including osteoarthritis, rheumatoid arthritis, septic arthritis, osteopenias such as osteoporosis, tumor metastasis (invasion and growth), periodontitis, gingivitis, corneal ulceration, dermal ulceration, gastric ulceration, and other diseases related to connective tissue degradation. Although there is a high incidence of these diseases in the developed world, there is no treatment that prevents the tissue damage that occurs. Considerable lines of scientific evidence indicate that uncontrolled connective matrix metalloproteinase (MMPs) activity is responsible for the damage, and as a consequence the inhibition of these enzymes has become the target for therapeutic intervention (see Matrisian, L. M., *Bases*, Vol. 14, pp 445-463, (1992); Emonard, H. et al., *Cellular and molecular Biology*, Vol. 36, pp 131-153, (1990); Docherty, A. J. P. et al., *Annals of the Rheumatic*, Vol. 49, pp 469-479, (1990)).

Hydroxamic acid derivatives are a class of known therapeutically active
25 MMPs inhibitors and there are numerous references in the art disclosing a variety
of hydroxamic acid derivatives. This invention provides α -amino sulfonyl
hydroxamic acids wherein the hydrogen on the amino nitrogen is unsubstituted and
wherein there is a side chain on the α -carbon position. The compounds of the
present invention have unexpected superior activity in inhibiting various enzymes
30 from the matrix metalloproteinase family, predominantly gelatinase, and hence are
useful for the treatment of matrix metallo endoproteinase diseases such as
osteoporosis, tumor metastasis (invasion and growth), periodontitis, gingivitis,
corneal ulceration, dermal ulceration, gastric ulceration, inflammation, and other
diseases related to connective tissue degradation.

INFORMATION DISCLOSURE

The following patent publications disclose sulfonamide hydroxamic acids as matrix metalloproteinase inhibitors:

European Patent Publication 0,606,046 A1 discloses arylsulfonamido-substituted hydroxamic acids useful as matrix metalloproteinase inhibitors.

International Publication No. WO 95/35275 and WO 95/35276 disclose hydroxamic acid and carboxylic acid derivatives useful as matrix metalloproteinase inhibitors.

International Publication No. WO 96/40101 A1 discloses new use of arylsulfonamido-substituted hydroxamic acids in the treatment of carcinoma and tumor angiogenesis.

International Publication No. WO 96/27583 A1 discloses arylsulfonamido N-hydroxamic acid derivatives of butyric acid useful as metalloproteinase inhibitors.

International Publication No. WO 96/33172 A1 discloses arylsulfonyl hydroxamic acid derivatives as matrix metalloproteinase and TNF inhibitors.

International Publication No. WO 97/20824 discloses benzenesulfonyl hydroxamic acids as metalloproteinase inhibitors.

International Publication No. WO 97/18194 A1 discloses cyclic and heterocyclic N-substituted α -iminohydroxamic and carboxylic acids as metalloproteinase inhibitors.


European Patent Publication 757984 A1 discloses hydroxamic acid derivatives as gelatinase inhibitors useful treating rheumatoid diseases.

European Patent Publication 757037 A2 discloses arylsulfonamidoamino acid derivatives useful as matrix metalloproteinase inhibitors.

MacPherson, L. J. at el. *J. Med. Chem.* Vol. 40, pp 2525-2523, (1997) discloses structure-activity relationships of a lead hydroxamic acids as matrix metalloproteinase inhibitors.

SUMMARY OF THE INVENTION

The present invention provides a compound of formula I



I

or pharmaceutical acceptable salts thereof wherein R₁ is isopropyl, 2-methylbut-2-yl, phenyl, benzyl, or 1H-indol-3-yl-methyl; and R₂ is *n*-octyl, phenyl, or phenyl substituted with methoxy, fluoro, or bromo.

The compounds of the present invention inhibit various enzymes from the
5 matrix metalloproteinase family, predominantly gelatinase, and hence are useful as preventatives and therapeutics for diseases related to connective tissue degradation.

DETAILED DESCRIPTION OF THE INVENTION

For the purpose of the present invention, the term "pharmaceutically
10 acceptable salts" refers to salts useful for administering the compounds of this invention and these include hydrochloride, hydrobromide, hydroiodide, trifluoroacetic acid, sulfate, phosphate, acetate, propionate, lactate, mesylate, maleate, malate, succinate, tartrate, citrate, 2-hydroxyethyl sulfonate, fumarate and the like. These salts may be in hydrated form. Some of the compounds of this invention may form
15 metal salts such as sodium, potassium, calcium and magnesium salts and these are embraced by the term "pharmaceutically acceptable salts".

The compounds of the present invention can be converted to their salts according to conventional methods.

The R₁ substituent is isopropyl, 2-methylbut-2yl, phenyl, benzyl, or 1H-indol-
20 3ylmethyl.

The R₂ substituent is *n*-octyl, phenyl, *p*-methoxyphenyl, *p*-fluorophenyl, or *p*-bromophenyl.

The compounds of formula I of this invention contain a chiral center at α -position of amino acids, as such there exist two enantiomers or a racemic mixture of
25 both. This invention relates to the compounds having *R*-configuration under the Cahn-Ingold-Prelog nomenclature system.

The compounds of this invention include the followings:


- a. N-Hydroxy-2(R)-[(4-methoxybenzenesulfonyl)amino]-3-(3-indolyl)-propanamide,
- 30 b. N-Hydroxy-2(R)-[(benzenesulfonyl)amino]-3-(3-indolyl)-propanamide,
- c. N-Hydroxy-2(R)-[(4-fluorobenzenesulfonyl)amino]-3-(3-indolyl)-propanamide,
- d. N-Hydroxy-2(R)-[(4-methoxybenzenesulfonyl)amino]-3-phenyl-propanamide,
- e. N-Hydroxy-2(R)-[(4-bromobenzenesulfonyl)amino]-3-(3-indolyl)-propanamide,
- f. N-Hydroxy-2(R)-[(*n*-octylsulfonyl)amino]-3-(3-indolyl)-propanamide,
- 35 g. N-Hydroxy-2(R)-[(4-methoxybenzenesulfonyl)amino]-2-phenyl acetamide,
- h. N-Hydroxy-2(R)-[(4-methoxybenzenesulfonyl)amino]-3-methyl butanamide,

- i. N-Hydroxy-2(R)-[(benzenesulfonyl)amino]-3-methyl butanamide, or
- j. N-Hydroxy-2(R)-[(4-methoxybenzenesulfonyl)amino]-3,3-dimethyl pentanamide.

The compounds of this invention can be prepared in accordance to the process
5 discussed below.

In CHART I, R_1 and R_2 are the groups as defined previously; R' is either hydrogen or a low alkyl or an (un)substituted phenyl. Structure 1 are sulfonylated in the presence of an appropriate base such as tertiary amine or pyridine to afford sulfonamide 2 or 3 directly. The reaction may be carried out in a solvent which does
10 not adversely affect the reaction (e.g. dichloromethane, diethylether, tetrahydrofuran or a mixture thereof). All of the starting amino acid acids or esters in the present invention are commercially available or can be prepared readily by one having ordinary skill in the art of organic chemistry, utilizing well known reactions. Saponification of compound 2 to carboxylate 3 occurs efficiently under either acidic
15 or basic conditions. Direct coupling of carboxylate 3 with hydroxylamine provides the desired hydroxamates 5.

CHART I



In this reaction, chromatography on silica gel is generally employed to purify the hydroxamates. Alternatively, carboxylate 3 is reacted with O-benzylhydroxylamine. This reaction provides a coupled product 4 which is more soluble in organic solvents, therefore, to allow easy isolation. Hydrogenolysis of the benzyl group by the methods well known to those skilled in the art afford compound 5.

The pharmaceutical compositions of this invention may be prepared by combining the compounds of formula I of this invention with a solid or liquid pharmaceutically acceptable carrier, and optionally, with pharmaceutically acceptable adjuvants and excipients employing standard and conventional techniques. Solid form compositions include powders, tablets, dispersible granules, capsules and suppositories. A solid carrier can be at least one substance which may also function

WO 98/17645

as a diluent, flavoring agent, solubilizer, lubricant, suspending agent, binder, tablet disintegrating agent, and encapsulating agent. Inert solid carriers include magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, cellulosic materials, low melting wax, cocoa butter, and the like.

5 Liquid form compositions include solutions, suspensions and emulsions. For example, there may be provided solutions of the compounds of this invention dissolved in water, water-propylene glycol, and water-polyethylene glycol systems, optionally containing conventional coloring agents, flavoring agents, stabilizers and thickening agents.

10 The pharmaceutical composition is provided by employing conventional techniques. Preferably the composition is in unit dosage form containing an effective amount of the active component, that is, the compounds of formula I according to this invention.

The quantity of active component, that is the compounds of formula I
15 according to this invention, in the pharmaceutical composition and unit dosage form thereof may be varied or adjusted widely depending upon the particular application method, the potency of the particular compound and the desired concentration. Generally, the quantity of active component will range between 0.5% to 90% by weight of the composition.


20 In therapeutic use for treating a patient, suffering from or susceptible to diseases involving connective tissue degradation, or inhibiting various enzymes from the matrix metalloproteinase family, including collagenase, stromelysin, and gelatinase, the compounds or pharmaceutical compositions thereof will be administered orally, parenterally and/or topically at a dosage to obtain and maintain
25 a concentration, that is, an amount, or blood-level of active component in the patient undergoing treatment which will be effective to inhibit such enzymes. Generally, an effective amount of the active compound will be in the range of about 0.1 to about 100 mg/kg. It is to be understood that the dosages may vary depending upon the requirements of the patient, the severity of connective tissue degradation being
30 treated, and the particular compounds being used. Also, it is to be understood that the initial dosage administered may be increased beyond the above upper level in order to rapidly achieve the desired blood-level or the initial dosage may be smaller than the optimum and the daily dosage may be progressively increased during the course of treatment depending on the particular situation. If desired, the daily dose
35 may also be divided into multiple doses for administration, e.g., two to four times per day.

The compounds of the present invention inhibit various enzymes from the matrix metalloproteinase family, predominantly gelatinase, and hence are useful for the treatment of matrix metalloproteinase diseases such as osteoarthritis, rheumatoid arthritis, septic arthritis, osteopenias such as osteoporosis, tumor metastasis (invasion and growth), periodontitis, gingivitis, corneal ulceration, dermal ulceration, gastric ulceration, various enzymes from the matrix metalloproteinase family, predominantly collagenase and other diseases related to connective tissue degradation. Such diseases and conditions are well known and readily diagnosed by physician of ordinary skill.

Pharmaceutical compositions for parenteral administration will generally contain a pharmaceutically acceptable amount of the compounds according to formula I as a soluble salt (acid addition salt or base salt) dissolved in a pharmaceutically acceptable liquid carrier such as, for example, water-for-injection and a suitably buffered isotonic solution having a pH of about 3.5-6. Suitable buffering agents include, for example, trisodium orthophosphate, sodium bicarbonate, sodium citrate, N-methylglucamine, L(+)-lysine and L(+)-arginine, to name a few. The compounds according to formula I generally will be dissolved in the carrier in an amount sufficient to provide a pharmaceutically acceptable injectable concentration in the range of about 1 mg/ml to about 400 mg/ml. The resulting liquid pharmaceutical composition will be administered so as to obtain the above-mentioned inhibitory effective amount of dosage. The compounds of formula I according to this invention are advantageously administered orally in solid and liquid dosage forms.

The compounds and their preparations of the present invention will be better understood in connection with the following examples, which are intended as an illustration of and not a limitation upon the scope of the invention.

EXAMPLE 1 Preparation of N-hydroxy-2(R)-[(4-methoxybenzenesulfonyl)-amino]-3-(3-indolyl)-propanamide



Step 1. Preparation of 2(R)-[(4-methoxybenzenesulfonyl)amino]-3-(3-indolyl)-propanoic acid methyl ester.

To a suspension of D-tryptophan methyl ester hydrochloride (4 mmol) in 40 mL of methylene chloride, magnetically stirred and cooled in an ice bath, under a nitrogen atmosphere, is added 8 mmol of NMM and 4 mmol of 4-methoxybenzene sulfonyl chloride. The mixture is allowed to come to ambient temperature overnight. It is diluted with ethyl acetate and washed twice with aqueous 10% HCl, then successively with water, 1 M sodium bicarbonate, and brine. The organic solution is dried over sodium sulfate and concentrated to afford 1.05 g (65%) of a white solid, m.p. 120-122 °C

Step 2. Preparation of 2(R)-[(4-methoxybenzenesulfonyl)amino]-3-(3-indolyl)-propanoic acid.

To 1.3 mmol of 2(R)-[(4-methoxybenzenesulfonyl)amino]-3-(3-indolyl)-propanoic acid methyl ester suspended in 15 mL of ethanol is added 5 mL of aqueous 2.5 M sodium hydroxide. The solid suspension dissolved, and is allowed to stir at ambient temperature overnight. The mixture is acidified with aqueous 10% HCl and extracted with ethyl acetate. The organic phase is washed with brine and dried over sodium sulfate. It is concentrated to afford 0.45 g (93%) of a white solid.


Step 3. Preparation of N-hydroxy-2(R)-[(4-methoxybenzenesulfonyl)amino]-3-(3-indolyl)-propanamide.

To 2(R)-[(4-methoxybenzenesulfonyl)amino]-3-(3-indolyl)-propanoic acid (0.67 mmol) in 6 mL of methylene chloride and 1 mL of DMF, cooled in an ice-bath, under a nitrogen atmosphere, is added 0.83 mmol of HOBT, 0.72 mmol of EDC, and 0.73 mmol of NMM. The mixture is allowed to stir for one hour. In a small test tube, 1 mmol of hydroxylamine hydrochloride and 0.9 mmol of NMM in 1 mL of DMF is agitated, forming a fine suspension which is then introduced into the carbodiimide reaction mixture. The mixture is allowed to come to ambient temperature overnight. It is diluted with 50 mL each of aqueous 10% HCl and ethyl acetate. The organic phase is washed with 10% HCl, twice with 1 M sodium bicarbonate, and brine. It is dried over sodium sulfate, concentrated, and chromatographed on silica gel, eluting with 30% acetone and 1% acetic acid in methylene chloride. Fractions containing the hydroxamic acid product are pooled and concentrated to afford 135 mg (52%) of a white solid, ¹H NMR (DMSO-d₆) δ 10.75, 10.6, 8.79, 7.91, 7.46, 7.24, 6.99, 6.97, 6.87, 6.82, 3.76, 3.72, 2.91, 2.64. MS (FAB) m/z 779, 390, 389, 329, 130. IR (mull) cm⁻¹ 3324, 2925, 1665, 1595, 1497, 1458, 1261, 1157. [α]_D = + 66° (0.62, ethanol).

EXAMPLE 2

Preparation of N-hydroxy-2(R)-[(benzenesulfonyl)amino]-3-(3-indolyl)-propanamide.

5




Following the general procedure outlined in EXAMPLE 1 (steps 1-3), and making non-critical variations but starting with benzenesulfonyl chloride, the title compound is obtained as a white solid.

^1H NMR ($\text{DMSO}-d_6$) δ 0.8, 10.6, 8.79, 8.13, 7.59-7.22, 7.03-6.86, 3.79, 2.93, 2.64
MS (EI) m/z : 359, 202, 157, 130. $[\alpha]_D = +38^\circ$ (1.75, ethanol).

15 EXAMPLE 3

Preparation of N-hydroxy-2(R)-[(4-fluorobenzenesulfonyl)amino]-3-(3-indolyl)-propanamide.

20




Following the general procedure outlined in EXAMPLE 1 (steps 1-3), and making non-critical variations but starting with 4-fluorobenzenesulfonyl chloride, the title compound is obtained.

^1H NMR (d_6 -DMSO) δ 10.8, 10.7, 8.90, 8.25, 7.56-7.6, 7.38, 7.31, 7.00-7.14, 6.96, 3.86, 2.76-2.10. MS (FAB) m/z : 378, 378, 147, 130, 73, 69, 58, 57, 55, 43, 41.

30 EXAMPLE 4

Preparation of N-hydroxy-2(R)-[(4-methoxybenzenesulfonyl)amino]-3-phenyl-propanamide.

35




WO 98/17645

Following the general procedure outlined in EXAMPLE 1 (steps 1-3), and making non-critical variations but starting with D-phenylalanine methyl ester hydrochloride, the title compound is obtained.

^1H NMR (d_6 -DMSO) δ 10.7, 8.88, 8.05, 7.56, 7.21-7.23, 7.09-7.11, 6.98, 3.86, 3.79-3.82, 2.56-2.85. MS (FAB) m/z : 351, 290, 236, 123, 75, 57.

EXAMPLE 5

Preparation of N-hydroxy-2(R)-[(4-bromobenzenesulfonyl)amino]-3-(3-indolyl)-propanamide.



Step 1. Preparation of 2(R)-[(4-bromobenzenesulfonyl)amino]-3-(3-indolyl)-propanoic acid methyl ester.


To 1 g (3.9 mmol) of D-tryptophan methyl ester hydrochloride in 12 mL of pyridine at ambient temperature is added 1 g (3.9 mmol) of 4-bromobenzene sulfonyl chloride. The yellow mixture is allowed to stir overnight. It is then poured into aqueous 10% HCl and extracted with several portions of ethyl acetate. The combined organic phases are washed with aqueous 10% HCl, water, and brine, and dried over sodium sulfate. The solution is decolorized with activated charcoal, and concentrated to 1.1 g (65%) of a white solid.

Step 2. Preparation of N-hydroxy-2(R)-[(4-bromobenzenesulfonyl)amino]-3-(3-indolyl)-propanamide.

Following the general procedure outlined in EXAMPLE 1 (steps 2 and 3), and making non-critical variations but starting with 2(R)-[(4-bromobenzenesulfonyl)amino]-3-(3-indolyl)-propanoic acid methyl ester, the title compound is obtained as an off-white solid. ^1H NMR ($\text{DMSO}-d_6$) δ 10.7, 8.86, 8.29, 7.41, 7.34, 7.28, 7.24, 7.03-6.98, 6.88, 3.78, 2.90, 2.70. IR (mull) cm^{-1} 2924, 1665, 1458, 1161, 741. MS (FAB) m/z 440, 439, 438, 437, 379, 377, 130. $[\alpha]_D = +61^\circ$ (0.7, methanol).

EXAMPLE 6

Preparation of N-hydroxy-2(R)-[(n-octylsulfonyl)amino]-3-(3-indolyl)-propanamide.



5


Following the general procedure outlined in EXAMPLE 7 (step 1) and making non-critical variations but starting with n-octylsulfonyl chloride, 2(R)-[(n-octylsulfonyl)amino]-3-(3-indolyl)-propanoic acid methyl ester is obtained as a colorless oil. ^1H NMR (CDCl_3) δ 8.28, 7.57, 7.35, 7.19, 7.12, 7.05, 5.0, 4.44, 3.72, 3.32, 3.24, 2.73, 1.65-1.50, 1.29-1.15, 0.88.

Following the general procedure outlined in EXAMPLE 1 (steps 2-3) and making non-critical variations but starting with 2(R)-[(n-octylsulfonyl)amino]-3-(3-indolyl)-propanoic acid methyl ester, the title compound is obtained as a glassy solid. ^1H NMR ($\text{DMSO}-d_6$) δ 10.9, 10.8, 9.0, 7.62, 7.54, 7.31, 7.17, 7.07, 7.0, 3.9, 3.05-2.9, 2.4, 1.3-0.9, 0.86. MS (EI) m/z : 395, 334, 174, 157, 130. $[\alpha]_D = +30^\circ$ (0.8, ethanol).

EXAMPLE 7

Preparation of N-hydroxy-2-[(4-methoxybenzenesulfonyl)amino]-2-phenyl acetamide.

20



25

Step 1. Preparation of 2(R)-[(4-methoxybenzenesulfonyl)amino]-2-phenyl acetic acid methyl ester.

To 6 mmol of (R)-phenyl glycine methyl ester (hydrochloride salt $[\alpha]_D = -111^\circ$ (1.34, aqueous 10% HCl)) in 10 mL of pyridine at ambient temperature is added 6.7 mmol of 4-methoxybenzene sulfonyl chloride. The yellow mixture is allowed to stir overnight. It is then poured into aqueous 10% HCl and extracted with several portions of ethyl acetate. The combined organic phases are washed with aqueous 10% HCl, water, and brine, and dried over sodium sulfate. The solution is concentrated to 2 g (quantitative yield) of a solid.

Step 2. Preparation of 2(R)-[(4-methoxybenzenesulfonyl)amino]-2-phenyl acetic acid.


WO 98/17645

To a suspension of 3 mmol of 2(R)-[(4-methoxybenzenesulfonyl)amino]-2-phenyl acetic acid methyl ester in 15 mL of ethanol is added 5 mL of aqueous 2.5 M sodium hydroxide. The solid suspension dissolved, and is allowed to stir, stoppered, at ambient temperature overnight. The resulting suspension is acidified with aqueous 10% HCl and extracted with ethyl acetate, adding brine to facilitate phase separation. The organic phase is washed with water and with brine and dried over sodium sulfate. It is concentrated to afford 0.95 g (quantitative yield) of a white solid.

Step 3. Preparation of N-hydroxy-2(R)-[(4-methoxybenzenesulfonyl)amino]-2-phenyl acetamide.

Following the general procedure outlined in EXAMPLE 1 (step 3), and making non-critical variations but starting with 2(R)-[(4-methoxybenzenesulfonyl)amino]-2-phenyl acetic acid, the title compound is obtained as a white solid. ^1H NMR (DMSO- d_6) δ 10.85, 8.91, 8.44, 7.57, 7.23-7.16, 6.90, 4.78, 3.76. IR (mull) cm^{-1} 3261, 2924, 1639, 1596, 1453, 1333, 1263, 1157. MS (FAB) m/z : 337, 276, 150. $[\alpha]_D = -4.4^\circ$ (0.87, ethanol). Calc'd for $\text{C}_{15}\text{H}_{16}\text{N}_2\text{O}_5\text{S}$: C, 53.6; H, 4.79; N, 8.33. Found: C, 53.55; H, 4.84; N, 8.25.

EXAMPLE 8 Preparation of N-hydroxy-2(R)-[(4-methoxybenzenesulfonyl)amino]-3-methyl butanamide.



Step 1. Preparation of N-benzyloxy-2(R)-[(4-methoxybenzenesulfonyl)amino]-3-methyl butanamide.


To a solution of N-(p-methoxybenzenesulfonyl)-D-valine (1.01 g, 3.53 mmol) in CH_2Cl_2 (25 mL) are added the following reagents in the order given: HOBT (477 mg, 3.53 mmol), 4-methylmorpholine (1.95 mL, 17.7 mmol), O-benzylhydroxylamine hydrochloride (1.69 mg, 10.6 mmol), and EDC (880 mg, 4.59 mmol). The resultant slurry is stirred overnight (20 h) at room temperature. The solvent is evaporated and the residual material purified by silica gel chromatography (50 g of SG, EtOAc) to give 1.89 g of an impure solid; the solid is reconstituted in EtOAc (150 mL) and washed with 1 N HCl (3 x 50 mL) and brine (50 mL). The organic layer is dried over anhydrous Na_2SO_4 , filtered, and concentrated to yield 1.11 g (79%) of product. The

material is recrystallized from hot EtOAc/hexane to yield 779 mg of the title compound, as a white solid. mp 156-158 °C; $[\alpha]_D^{25} +11^\circ$ (c 1.01, CHCl₃); IR (mineral oil) 3252, 1657, 1596, 1502, 1445, 1354, 1328, 1306, 1266, 1165, 1160, 1146, 1096, 748, 671 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 8.56 , 7.77 , 7.36, 6.96, 5.22, 4.65-4.80, 3.84, 3.30-3.40, 1.90-2.05 , 0.70-0.90.

Step 2 . Preparation of N-hydroxy-2(R)-[(4-methoxybenzenesulfonyl)amino]-3-methyl butanamide.

A solution of N-benzyloxy-2(R)-[(4-methoxybenzenesulfonyl)amino]-3-methyl butanamide, the product of step 1, in 50% MeOH/EtOAc (30 mL) is degassed with N₂ and treated with Pearlman's catalyst (160 mg). The atmosphere is replaced with H₂ via a balloon. After 3 hours, the reaction mixture is filtered through celite, washing the residual cake with excess MeOH and EtOAc. The filtrate is concentrated to give 620 mg of a white solid. The material is crystallized from hot MeOH/CH₂Cl₂ and hexane to give the title compound (345 mg , 56%) as a white, crystalline solid (mp 166-168 °C). The mother liquor also yielded 251 mg (for a total yield of 97%) of additional product. $[\alpha]_D^{25} -4^\circ$ (c 0.93, DMSO); IR (mineral oil) 3269, 1634, 1599, 1539, 1497, 1337, 1310, 1261, 1162, 1096, 1025, 845, 804, 674, 629 cm⁻¹; ¹H NMR (300 MHz, DMSO-d₆) δ 10.51, 8.81, 7.65-7.85, 7.69, 7.05, 3.83, 3.20-3.30, 1.65-1.80, 0.60-0.90; MS (EI) *m/z* 302 (M⁺), 259, 242, 171, 107, 92, 76; Anal: C, 47.76; H, 6.30; N, 9.33; S, 10.27.

EXAMPLE 9 Preparation of N-hydroxy-2(R)-[(benzenesulfonyl)amino]-3-methyl butanamide.



Step 1. Preparation of (R)-[2-methyl-1-[(phenylmethoxy)amino]-carbonyl]propyl]-carbamic acid, 1,1-dimethylethyl ester.

CDI (2.45 g, 15.1 mmol) is added to a solution of N-(tert-butoxycarbonyl)-D-valine (3.28 g, 15.1 mmol) and CH₂Cl₂ (60 mL). The solution is stirred for 1 hour at room temperature. Diisopropylethylamine (2.90 mL, 16.6 mmol) and O-benzylhydroxylamine (2.64 g, 16.5 mmol) are added and the solution stirred for 16 hours at room temperature. The solution is concentrated, diluted with EtOAc, and

WO 98/17645

washed with 5% HCl (2 x 50 mL), NaHCO₃ (50 mL) and brine (50 mL). The organic solution is dried (MgSO₄), filtered, and concentrated to give 4.42 g (91%) of the title compound as a white solid which is carried on crude.

Step 2. Preparation of D-2-amino-N-(benzyloxy-3-methyl-butylamide).

5 TFA (8.0 mL) is added to a solution of (R)-[2-methyl-1-[(phenylmethoxy)-amino]carbonyl] propyl]-carbamic acid, 1,1-dimethylethyl ester, the product of step 1, (1.00 g, 3.10 mmol) and CH₂Cl₂ (10.0 mL) at 0 °C. The solution is stirred at 0 °C for 1 hour and is concentrated. Basic workup (CH₂Cl₂, NaHCO₃, MgSO₄) gives 665 mg (96%) of the title compound as a white solid which is carried on crude.

10 Step 3. Preparation of N-benzyloxy-2(R)-[(benzenesulfonyl)amino]-3-methyl butanamide.

Benzenesulfonyl chloride (280 µL, 2.19 mmol) is added to a cold (0 °C) solution of D-2-amino-N-(benzyloxy)-3-methyl-butyramide, the product of step 2, (442 mg, 1.99 mmol) and Hunig's base (380 µL, 2.18 mmol) in CH₂Cl₂ (20 mL).

15 After 1 hour at 0 °C, the solution is allowed to warm to room temperature overnight. The reaction mixture is partitioned between CH₂Cl₂ and sat. NaHCO₃. The organic layer is dried (MgSO₄), filtered, and concentrated. The desired material is crystallized from CH₂Cl₂/MeOH/pentane to give 481 mg (67%) of the title compound as a crystalline solid. mp 163-165 °C.

20 Step 4. Preparation of N-hydroxy-2(R)-[(benzenesulfonyl)amino]-3-methyl butanamide.


A suspension of N-benzyloxy-2(R)-[(benzenesulfonyl)amino]-3-methyl butanamide (100 mg, 0.280 mmol) and Pearlman's catalyst (25 mg) in MeOH (10 mL) is hydrogenated at 1 atmosphere for 3 hours. The mixture is filtered through celite, and the filtrate is concentrated. The desired material is crystallized from EtOAc/pentane to give 61 mg (80%) of the title compound as a crystalline solid. mp 154-155 °C; IR (mineral oil) 3268, 2925, 2954, 2881, 2855, 1636, 1449, 1336, 1165, 694 cm⁻¹; ¹H NMR (300 MHz, DMSO-*d*₆) δ 10.53, 8.83, 7.98, 7.76, 7.50-7.65, 3.28, 1.65-1.85, 0.74, 0.70; MS (EI) *m/z* 272; Analysis: C, 48.37; H, 6.07; N, 10.14.

30

EXAMPLE 10

Preparation of N-Hydroxy-2-[(4-methoxybenzenesulfonyl)-amino]-3,3-dimethyl pentanamide.

35



5

Step 1. Preparation of 2-[(4-methoxybenzenesulfonyl)amino]-3,3-dimethyl pentanoic acid.

2-amino-3,3-dimethylpentanoic acid hydrochloride (1.63 g), (JCS Chem. Comm. 11, 830-1992) is mixed with diisopropylethylamine (3.6 g, 3.1 equivalent) in dry THF (100 mL). The 4-methoxybenzenesulfonyl chloride (1.86 g, 1 equivalent) is added as a solid. DMF (25 mL) is added to the resulting suspension. After stirring overnight at room temperature the solvents are removed under vacuum. The residue is partitioned between ethyl acetate (200 mL) and 1N HCl (100 mL). The organic phase is separated and the aqueous phase extracted with several more portions of ethyl acetate. The combined organic phases are washed with 1N HCl and then brine. The organic phase are dried (MgSO₄), filtered, and concentrated under reduced pressure. The resulting crude product is loaded onto a reverse phase column as a DMSO solution. The column is eluted using a step gradient (10% increments), starting with 30% acetonitrile/water and ending at 90% acetonitrile/water. Combination of the appropriate fractions gave 116 mg of the title compound.

Step 2. Preparation of N-hydroxy-2-[(4-methoxybenzenesulfonyl)amino]-3,3-dimethyl pentanamide.

The product of step 1 (0.116 g), EDC hydrochloride (0.076 g, 1.1 equivalent), hydroxybenzotriazole (0.054 g, 1.1 equivalent), and diisopropylethylamine (0.143 g, 3 equivalent) are mixed with dichloromethane (20 mL) to give a homogeneous solution. Hydroxylamine hydrochloride (0.051 g, 2 equivalent) is added as a solid. DMF (3 mL) is added to solubilize the hydroxylamine hydrochloride. The reaction is allowed to stir at room temperature overnight. The solvents are then removed under reduced pressure and the residue partitioned between ethyl acetate and 1N HCl. The organic phase is separated and the aqueous phase re-extracted with ethyl acetate. The combined organic phases are dried (MgSO₄), filtered, and concentrated under reduced pressure. The crude product is loaded onto a reverse phase column as a DMSO solution. The column is eluted with a step gradient, starting at 20% acetonitrile/water, and raised in 10% increments. The appropriate fractions are combined to give 22 mg of the title compound.

^1H NMR (DMSO, 300 MHz): 10.4, 7.69-7.65, 7.43, 7.01-6.98, 3.80, 1.20-1.17, 0.77, 0.75, 0.69.

EXAMPLE 11 Biological Activity Test

5 Inhibitory activity is evaluated in one or more of the MMP enzymes (stromelysin, gelatinase, and collagenase) in vitro using particle concentration fluorescence assay. An inhibitor binds to MMP enzymes which prevents the degradation of a substrate by stromelysin, gelatinase, or collagenase. The substrate has attached to it a fluorescein and a biotin moiety. The intact substrate then binds
10 to an avidin-coated particle via the biotin moiety. Once the particle is washed and dried, a fluorescent signal is generated since the fluorescent group is attached to the particle. Without an inhibitor present, the substrate is degraded by MMP enzymes and the fluorescein group is removed, therefore, no fluorescent signal can be detected. Testing compounds are dissolved in DMSO to the desired concentration,
15 then the solutions are diluted to 1:5 with MMP buffer (50 mM Tris-HCl, pH 7.5; 150 mM NaCl; 0.02% NaN₃). Serial two-fold dilutions of each compound are prepared. A concentrated, activated enzyme solution is transferred into each plate of the testing compounds, and the mixture is incubated at room temperature for 15 minutes. Thawed MMP substrate is then added into all plates, and the plates are
20 incubated in the dark for 1-3 hours at room temperature. At this point, the substrate mixture is mixed with 0.1% avidin-coated polystyrene particles. After 15 minutes, the fluorescence values are measured following filtration and washing of the beads. K_i values are then calculated. Inhibitory data for the compounds of this invention are shown in TABLE 1. Compounds with lower K_i values are expected to
25 be more effective as MMP inhibitors. It is expected that a compound with a K_i less than 15 μM against stromelysin, collagenase, or gelatinase will display therapeutic effects in connective tissue disorders.

TABLE 1


MMP Inhibition Constants (K_i , μM) of the Compounds of the Invention

5	Example No.	Gelatinase K_i (μM)	Example No.	Gelatinase K_i (μM)
	1	0.00781	2	0.0142
	3	0.079	4	0.00723
	5	0.0026	6	0.0121
	7	0.0033	8	0.0091
10	9	0.082	10	0.0098

CLAIMS

We claim:

1. A compound of formula I



I

or pharmaceutical acceptable salts thereof wherein

R₁ is

- a) isopropyl,
- b) 2-methylbut-2-yl
- c) phenyl,
- d) benzyl, or
- e) 1H-indol-3-ylmethyl; and

R₂ is

- a) *n*-octyl,
- b) phenyl, or
- c) phenyl substituted with methoxy, fluoro, or bromo.

2. A compound of claim 1 wherein R₁ is selected from the group consisting of isopropyl, 2-methylbut-2-yl, 1H-indol-3-ylmethyl, phenyl, and benzyl.

3. A compound of claim 1 wherein R₂ is selected from the group consisting of *n*-octyl, phenyl, *p*-methoxyphenyl, *p*-fluorophenyl, or *p*-bromophenyl.

4. A compound of claim 1 which is

- a. N-Hydroxy-2(R)-[(4-methoxybenzenesulfonyl) amino]-3-(3-indolyl)-propanamide,
- b. N-Hydroxy-2(R)-[(benzenesulfonyl)amino]-3-(3-indolyl)-propanamide,
- c. N-Hydroxy-2(R)-[(4-fluorobenzenesulfonyl)amino]-3-(3-indolyl)-propanamide,
- d. N-Hydroxy-2(R)-[(4-methoxybenzenesulfonyl)amino]-3-phenyl-propanamide,
- e. N-Hydroxy-2(R)-[(4-bromobenzenesulfonyl)amino]-3-(3-indolyl)-propanamide,
- f. N-Hydroxy-2(R)-[(*n*-octylsulfonyl)amino]-3-(3-indolyl)-propanamide,

- g. N-Hydroxy-2(R)-[(4-methoxybenzenesulfonyl)amino]-2-phenyl acetamide,
- h. N-Hydroxy-2(R)-[(4-methoxybenzenesulfonyl)amino]-3-methyl butanamide,
- i. N-Hydroxy-2(R)-[(benzenesulfonyl)amino]-3-methyl butanamide, or
- j. N-Hydroxy-2(R)-[(4-methoxybenzenesulfonyl)amino]-3,3-dimethyl
5 pentanamide.

5. A method of inhibiting excess matrix metalloproteinase which comprises administering to a patient in need thereof an effective amount of a compound of claim 1.

10

6. A method of claim 5 wherein matrix metalloproteinases comprises stromelysin, collagenase, and gelatinase.

7. A method of treating a human, suffering from or susceptible to a diseases
15 involving connective tissue degradation which comprises administering to a patient in need thereof an effective amount of a compound of claim 1.

8. A method of 7 wherein the diseases related to connective tissue degradation is osteoarthritis, rheumatoid arthritis, septic arthritis, osteopenias such as
20 osteoporosis, tumor metastasis (invasion and growth), periodontitis, gingivitis, corneal ulceration, dermal ulceration, or gastric ulceration.

9. The method of claim 5 wherein the effective amount of the compound of claim 1 is administered orally, parenterally, or topically in a pharmaceutical composition.

25

10. The method of claim 7 wherein the effective amount of the compound of claim 1 is administered orally, parenterally, or topically in a pharmaceutical composition.

11. The method of claim 5 or 7 wherein said compound is administered in an
30 amount of from about 0.1 to about 100 mg/kg of body weight/day.

12. A pharmaceutical composition which comprises an amount of the compound of claim 1 effective to inhibit excess matrix metalloproteinase and a pharmaceutically acceptable carrier.

35

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 97/18235

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07D209/20 A61K31/40 C07C311/29 C07C311/19

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07D A61K C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 606 046 A (CIBA-GEIGY AG) 13 July 1994 cited in the application see claims ---	1,12
A	WO 95 35276 A (BRITISH BIOTECH PHARMACEUTICALS LTD) 28 December 1995 cited in the application see examples 5,11,14,15 ---	1,12
P,A	EP 0 757 984 A (ONO PHARMACEUTICAL CO.,LTD.) 12 February 1997 * page 11, Table 1, examples 1,3,7 * ---	1,12
P,X	WO 97 27174 A (SHIONOGI & CO.,LTD.) 31 July 1997 * page 38-41: examples 9,12,26 ; page 53, example 72 * -----	1-4,12

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

17 February 1998

Date of mailing of the international search report

05.03.98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Van Bijlen, H

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 97/ 18235

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 5-11
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim(s) 5-11
is(are) directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 97/18235

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 606046 A	13-07-94	US 5455258 A	03-10-95
		AT 159012 T	15-10-97
		AU 684255 B	11-12-97
		AU 5265593 A	04-05-95
		CA 2112779 A	07-07-94
		DE 69314456 D	13-11-97
		FI 940012 A	07-07-94
		HU 70536 A	30-10-95
		JP 6256293 A	13-09-94
		MX 9400276 A	29-07-94
		NO 940038 A,B,	07-07-94
		NZ 250517 A	26-10-95
		US 5506242 A	09-04-96
		US 5552419 A	03-09-96
		US 5646167 A	08-07-97
		US 5672615 A	30-09-97
		ZA 9400048 A	11-08-94
WO 9535276 A	28-12-95	AU 2746595 A	15-01-96
		AU 2746695 A	15-01-96
		CA 2193691 A	28-12-95
		CA 2193692 A	28-12-95
		EP 0766664 A	09-04-97
		EP 0766665 A	09-04-97
		FI 965153 A	20-12-96
		WO 9535275 A	28-12-95
		GB 2303850 A	05-03-97
		GB 2303629 A	26-02-97
EP 757984 A	12-02-97	NO 965515 A	20-02-97
		JP 9104672 A	22-04-97
WO 9727174 A	31-07-97	AU 1319597 A	20-08-97